

PROJECTED CHANGES

| and the same of th |
|--|
| |
| |





| Decrease |
|----------------|
| in cold nights |
| per year |

| Increase |
|----------------|
| in severe fire |
| weather days |
| per year |

| Low-emissions scenario | | |
|------------------------|------|--|
| 2050 | 2090 | |
| +1.1°C | +1.2 | |
| +13.0 | +13. | |
| -12.3 | -13. | |
| +2.6 | +3.1 | |


| | nissions nario | | emissions nario |
|--------|-------------------|--------|--------------------|
| 2050 | 2090 | 2050 | 2090 |
| +1.1°C | +1.2°C | +1.5°C | +2.5°C |
| +13.0 | +13.8 | +16.1 | +28.3 |
| -12.3 | -13.5 | -15.5 | -24.7 |
| +2.6 | +3.1 | +2.8 | +6.1 |
| | | | |


| Medium-emissions scenario | High-emissions scenario |
|---------------------------|-------------------------|
| 2050 2090 | 2050 2090 |
| +1.5°C +2.5°C | +1.9°C +3.7°C |
| +16.1 +28.3 | +20.8 +41.9 |
| -15.5 -24.7 | -18.7 -31. 7 |
| +2.8 +6.1 | +4.8 +8.0 |

REGIONAL IMPACTS











Viticulture

Data is based on NARCliM2.0 projections for SSP1-2.6 (low-emissions), SSP2-4.5 (medium-emissions) and SSP3-7.0 (high-emissions) and is presented relative to the baseline period of 1990-2009. Values presented are averages across the NARCliM2.0 model ensemble, and do not represent the full range of plausible climate futures. Climate change impacts are used to highlight how the region is likely to be affected by climate change, and impacts are not limited to the examples provided.

Inland wetlands