

NARClIM

South East and Tablelands

Climate Change Snapshot




environment.nsw.gov.au



Photo caption:

The Emu in the Sky is an Aboriginal constellation that is made up of the dark clouds of the Milky Way. With the movement of the Earth, the position of the Emu in the Sky changes throughout the year. Aboriginal people in some nations across NSW and Australia relate the position of the Emu in the Sky to the breeding behaviour of the emu on the land. Cultural astronomy teaches us about the relationship between the sky and land; and that we are all interconnected.



Published by:

Department of Climate Change,
Energy, the Environment and Water
Locked Bag 5022, Parramatta
NSW 2124

T +61 2 9995 5000 (switchboard)
T 1300 361 967 (Environment and
Heritage enquiries)

TTY users: phone 133 677
then ask for 1300 361 967

Speak and listen users: phone
1300 555 727 then ask for
1300 361 967

E info@environment.nsw.gov.au

W www.environment.nsw.gov.au

ISBN 978-1-923516-82-3
Published: November 2025

Acknowledgement of Country

The NSW Government acknowledges First Nations people as the first Australian people and the traditional owners and custodians of the country's lands and water. Australia's First Nations people have lived in NSW for over 60,000 years and have significant spiritual, cultural and economic connections with its lands, waters, seas and skies.

The South East and Tablelands region encompasses the traditional lands of the Wiradjuri, Ngunnawal, Ngarigo, Tharawal, Gundungurra, Dharug, Yuin and Bidwell peoples.

They are the first astronomers and scientists who have been listening, reading and understanding natural processes and caring for Country for generations.

We pay respects to Elders past and present and acknowledge the significance of their traditional knowledge in adapting to changes in climate over tens of thousands of years.

We recognise the importance of their cultural knowledge and guidance at this pivotal moment in time.



About this snapshot

The New South Wales (NSW) and Australian Regional Climate Modelling (NARClIM) project delivers high-resolution climate change projections for NSW and south-east Australia.

This snapshot provides the latest NARClIM2.0 climate projections for the South East and Tablelands under low, medium and high emissions scenarios for the middle of the century (2050) and end of the century (2090). It includes projections for key climate variables including temperature, average rainfall, hot days (days $\geq 35^{\circ}\text{C}$), cold nights ($< 2^{\circ}\text{C}$), and severe fire weather days (Forest Fire Danger Index > 50). The projections help illustrate potential climate changes and their impacts, as well as associated climate risks.

NSW is already experiencing climate change. This document provides local-scale climate modelling insights to help the South East and Tablelands communities understand and plan for the impacts of climate change on their infrastructure, environment and way of life; and to support informed planning, risk assessment and action.

This snapshot offers a high-level overview, with more detailed data available through the [AdaptNSW Interactive Map](#), [Climate Data Portal](#) and [AdaptNSW](#).

How to use this snapshot

While there are several different ways to engage with the information in this snapshot, here are some key things to consider:

- **Explore each climate variable across scenarios** – review projections under low (SSP1–2.6), medium (SSP2–4.5), and high (SSP3–7.0) emissions scenarios to understand how climate risk differs depending on emissions pathways (Shared Socioeconomic Pathways, SSPs).
- **Compare scenario-based changes over time** – examine how each climate variable responds to different emissions scenarios for the middle of the century (2050) and the end of the century (2090) to understand how risks may evolve.
- **Identify where projections of climate variables align or diverge** – look for patterns across emissions scenarios and timeframes to see where risks remain consistent and where they escalate or diverge significantly.

Time periods in this snapshot

The projections for each time period represent averaged data across all climate models used for NARClIM for a 20-year period:

- **Baseline period: baseline** → The modelled average for each climate variable from 1990–2009, used for comparison with future projections.
- **Middle of the century: ‘2050’ projection** → The projected average for each climate variable for 2040–2059.
- **End of the century: ‘2090’ projection** → The projected average for each climate variable for 2080–2099.



About this snapshot

NARClIM climate projections

NARClIM is NSW's regional climate modelling project. NARClIM combines carefully selected global and regional climate models through a process known as dynamical downscaling, to generate detailed, locally relevant climate projections. These simulate a range of plausible future climates, helping to inform climate risk assessments and support planning at local and regional levels.

Launched in 2024, NARClIM2.0 provides nation-leading climate model data that span the range of plausible future changes in climate. It offers:

- climate projections out to the year 2100, and simulations of the past climate from 1951 to 2014
- 4-km scale projections for south-east Australia
- 20-km scale projections for the broader Australasian region
- projections for key climate variables and extremes.

There is more information [About NARClIM](#), as well as specific information on [Downscaling in NARClIM](#) and [Global and regional climate models used by NARClIM](#) at AdaptNSW.

Methods and uncertainty

To help address future uncertainty, NARClIM2.0 is built on a selection of emissions scenarios, global climate models and regional climate models that, together, capture a range of climates that could occur. This is referred to as the NARClIM model ensemble. The NARClIM2.0 model ensemble is made up of different combinations of 5 global climate models and 2 regional climate models, giving 10 model combinations in total.

The data presented in this snapshot is generally an average for different 20-year time periods (e.g. the 2050 projection is the average for the 2040–2059 time period). Time series data are presented as annual averages. Combining multiple models through averaging and other statistical methods produces better projections by providing a comprehensive representation of possible future climate scenarios.

To ensure that NARClIM models adequately simulate regional climate, scientists use them to simulate the past climate and compare the results with actual observations. Outputs undergo rigorous quality control and scientific technical peer review.

There is more information on [The NARClIM modelling methodology](#) and [NARClIM data processing, testing and validation](#) at AdaptNSW.

Mental health support

Climate change information can be distressing for some readers, with many Australians of all ages experiencing significant eco-anxiety. For supporting information, please visit the [Black Dog Institute](#) or [Australian Psychological Society](#) or speak with your local healthcare provider.

About this snapshot

Shared Socioeconomic Pathways

NARClIM2.0 uses Shared Socioeconomic Pathways (SSPs), which are the most recent emissions scenarios adopted in [Coupled Model Intercomparison Project Phase 6 \(CMIP6\)](#) models and used in the Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report.

The SSPs are a type of storyline-based emission scenario that estimate the world's future emissions and how these will affect the climate. SSPs outline different global development trajectories based on factors such as population, economic growth, education, urbanisation and land use, and technological advancement.¹ By analysing SSPs, we can better understand the long-term consequences of today's decisions and determine if we are heading toward higher-risk scenarios.²


For more information on emissions scenarios visit [Emissions scenarios used by NARClIM](#) on AdaptNSW and [Summary for policymakers report](#) by the IPCC.


Why do we use 3 SSPs?

The future is uncertain. There are many plausible futures on the horizon, and the one we reach depends on the path we take to get there. NARClIM provides projections for 3 SSPs (low-, medium- and high-emissions), each representing a distinct future with varying levels of climate risk.

Considering a range of SSPs and understanding where these scenarios align or diverge – in both the middle of the century (2050) and the end of the century (2090) – helps inform better planning and decision-making. NARClIM data highlights just how stark the differences between futures can be.

For more information about how to integrate this information into your risk assessments see [Climate risk ready guide](#) and [Limitations and appropriate use](#) on AdaptNSW.





About this snapshot

Understanding the baseline period

To assess future climate projections, a climate baseline is used. This is a reference point which future change is relative to. In this snapshot, the baseline is the 20-year period from 1990 to 2009. This period is termed the baseline period to represent the average climate across those 2 decades.

A 20-year baseline averages out natural climate variability and avoids misleading comparisons with unusually hot, cold, wet or dry years. Using a fixed reference point prevents issues that may arise from using shifting reference points to compare future change against.

Climate during the baseline period is described in 2 ways in this snapshot:

- **Historical model:** The NARClIM2.0 simulation of past climate conditions.
- **Observed:** What was actually measured using weather station data during this period.

These 2 values are similar but not the same. Climate models aim to capture long-term patterns and trends, rather than matching observations perfectly. Observed values give context for comparison of the historical model with what it was in reality.

Looking backwards from the baseline

Before the baseline period, +0.84°C of observed warming had already occurred across NSW and the Australian Capital Territory (ACT) since records began. This is the difference between the 20-year average temperature of the 2 periods centred on 1920 (1910–1929) and 2000 (1990–2009). The Bureau of Meteorology’s national climate records for temperature begin in 1910, making 1910–1929 the first available 20-year average for comparison with the baseline.

Consider the following when incorporating past warming into future projections:

- Warming before the baseline (+0.84°C) is not included in projections of future change.
- Warming after the baseline period is already included in future projections and should not be added again.

Looking forwards from the baseline

Use the historical model values in Table 1 as the baseline when interpreting both middle of the century and end of the century climate projections in this snapshot.

By comparing future projections to the historical model baseline values, we ensure the projected changes reflect genuine shifts, instead of also including the small differences between the modelled and observed data.

Table 1. Baseline climate for the South East and Tablelands

	Average temperature	Average maximum temperature	Average minimum temperature	Hot days	Cold nights	Rainfall	Severe fire weather days
Observed	13.0°C	19.3°C	6.6°C	4.9 days	86.8 days	723 mm	0.5 days
Historical model	12.7°C	18.3°C	8.0°C	4.4 days	88.9 days	743 mm	1.2 days

Table 1 outlines the annual average values for the baseline period in this snapshot. All observed data is calculated from Bureau of Meteorology products. Long-term temperature change data is from the long-term temperature record.³ Observed information and data in graphs come from Australian Gridded Climate Data (AGCD).⁴

Climate of the South East and Tablelands

The climate of the South East and Tablelands underpins a diverse array of important lifestyles, industries and natural ecosystems. A stable climate is critical to support a range of values in the South East and Tablelands, including our unique biodiversity, recreational activities and food systems.

The South East and Tablelands region encompasses the traditional lands of the Wiradjuri, Ngunnawal, Ngarigo, Tharawal, Gundungurra, Dharug, Yuin and Bidwell peoples.

The South East and Tablelands region is in the south-east corner of NSW, and includes cities, towns and settlements surrounding the ACT, including Batemans Bay, Bega, Cooma, Goulburn, Moss Vale, Queanbeyan, Yass and Young. The South East and Tablelands region offers snow, surf and rural living, with connections to Canberra and Sydney. The region is known for its beautiful beaches, large coastal embayments and Australia's highest summits, including Mount Kosciuszko. It also contains the headwaters of some of Australia's most famous rivers including the Lachlan, Murray, Murrumbidgee and Snowy.

The topography of the South East and Tablelands region results in a large range of climates. It is relatively wet close to the coast and Snowy Mountains, and drier inland. It is hot in summer in the northern inland areas and very cold in winter in the Snowy Mountains. Milder conditions are found along the coast, with cooler temperatures in summer and warmer temperatures in winter. The Southern Tablelands and south-eastern slopes contain many temperate grassland, woodland, wet forest and alpine ecosystems, and plant and animal species not found elsewhere in NSW.

People aged 65 and over make up 23.3% of the population, (notably higher than the NSW and ACT average of 17.4%), while those aged 0–14 years represent 17.2% and the working-age population (15–64 years) represent 59.9% of the region's total.⁵


The South East and Tablelands region supports a diverse range of industries that are vital for NSW's economy. The largest industries in

the region, by employment, are public administration and safety (12.7%), health care and social assistance (12.3%), construction (10.7%), retail trade (8.7%) and education and training (8.0%).⁵


The region's climate has provided the foundation for many of the region's current social, economic and ecological systems. These systems will be impacted by increased temperatures, more hot days, fewer cold nights, greater fire danger and higher rainfall variability.

The following pages outline the projected changes in these key climate variables across the South East and Tablelands region.





PROJECTED CHANGES



Increase in average temperature

Decrease in cold nights per year

Decrease in average winter rainfall

Increase in severe fire weather days per year

Low-emissions scenario		Medium-emissions scenario		High-emissions scenario	
2050	2090	2050	2090	2050	2090
+1.1°C	+1.2°C	+1.4°C	+2.5°C	+1.8°C	+3.6°C
-16.4	-19.8	-23.2	-39.3	-28.0	-55.2
-9.7%	-8.3%	-12.1%	-28.3%	-17.2%	-23.7%
+0.5	+0.7	+0.6	+1.7	+1.2	+2.4

REGIONAL IMPACTS


Alpine ecosystems
Decrease in cold nights

Isolated coastal towns
Increased severe fire weather

Agriculture
Increased extreme heat

River systems
Changes to rainfall

Data is based on NARClIM2.0 projections for SSP1–2.6 (low-emissions), SSP2–4.5 (medium-emissions) and SSP3–7.0 (high-emissions) and is presented relative to the baseline period of 1990–2009. Values presented are averages across the NARClIM2.0 model ensemble, and do not represent the full range of plausible climate futures. Regional climate change impacts are used to highlight how the region is likely to be affected by climate change, and impacts are not limited to the examples provided.



Temperature

In NSW, 8 of the 10 warmest years on record since 1910 have occurred since 2013.



3.6°C

rise in average temperature across the South East and Tablelands by 2090 under a high-emissions scenario.

Temperatures are projected to be higher by 2050 under a high-emissions scenario than by 2090 under a low-emissions scenario.

The South East and Tablelands is getting warmer

Temperature is the most robust indicator of climate change. The warmest year on record for both mean temperature and maximum temperature in the South East and Tablelands was 2019, when the average temperature was 0.9°C above the 1990–2009 baseline average.⁴

Projections

Across the South East and Tablelands region, average temperatures will increase throughout this century (Figure 1).

Under a low-emissions scenario, the average temperature increase across the region is projected to be 0.1°C between 2050 and 2090 (Table 2). However, major temperature increases of 1.1°C under a medium-emissions scenario and 1.8°C under a high-emissions scenario are expected during the same period. Notably, the temperature projections for 2050 under a medium-emissions scenario and a high-emissions scenario are expected to exceed the projections for 2090 under a low-emissions scenario.

Table 2 and Figure 1 provide more information on how the projections differ across the 3 scenarios, and Figure 2 provides information on regional differences by 2090 across the 3 scenarios.





Temperature

Table 2. Projected annual average temperature increase – South East and Tablelands

2050


	Low-emissions	Medium-emissions	High-emissions
Temperature	1.1°C (0.5°C to 1.5°C)	1.4°C (0.8°C to 1.9°C)	1.8°C (0.8°C to 2.6°C)
Maximum temperature	1.2°C (0.5°C to 1.7°C)	1.5°C (1.0°C to 2.0°C)	2.0°C (0.9°C to 2.9°C)
Minimum temperature	1.0°C (0.5–1.4°C)	1.4°C (0.8°C to 1.9°C)	1.7°C (0.8°C to 2.4°C)

2090

	Low-emissions	Medium-emissions	High-emissions
Temperature	1.2°C (0.5°C to 1.8°C)	2.5°C (1.7°C to 3.6°C)	3.6°C (2.3°C to 5.1°C)
Maximum temperature	1.3°C (0.5°C to 2.1°C)	2.7°C (1.8°C to 3.9°C)	3.8°C (2.5°C to 5.4°C)
Minimum temperature	1.1°C (0.6°C to 1.6°C)	2.3°C (1.7°C to 3.3°C)	3.5°C (2.3°C to 4.9°C)

The bold number is the ensemble average for the period. Underneath the average is the ensemble range. Temperature increases are additional to the historical model baselines of 12.7°C for average temperature, 18.3°C for average maximum temperature and 8.0°C for average minimum temperature.


Figure 1. Historical and projected average temperature change – South East and Tablelands






Temperature

Figure 2. Projected change in average temperature by 2090 for the South East and Tablelands





Hot days

Changes to temperature extremes often have more pronounced impacts than changes in average temperatures.

Hot days will become more frequent

Prolonged hot days, where maximum daily temperatures are equal to or above 35°C, increase the incidence of illness and death – particularly among vulnerable people. Seasonal changes in the number of hot days could have significant impacts on bushfire danger, infrastructure and native species.

The number of hot days varies widely across the South East and Tablelands region. During the baseline period, areas along the coast and higher-elevation alpine areas had on average less than 1 hot day per year. Inland areas, such as Moss Vale, Goulburn and Queanbeyan, had on average fewer than 5 hot days per year, while western areas of the region near Young had on average fewer than 20 hot days per year.

Projections

Across the South East and Tablelands, the average number of hot days per year will increase throughout this century (Figure 3).

The number of hot days will increase for the region by 2050 for all emissions scenarios, with an even greater increase by 2090 under a medium-emissions scenario and a high-emissions scenario (Table 3). The number of hot days is projected to increase across spring, summer and autumn, with the largest increase in summer.

Under a low-emissions scenario, there is a minimal increase of less than

5x

By 2090, the South East and Tablelands could experience nearly 5 times the number of hot days per year under a high-emissions scenario.

Higher maximum temperatures affect health through heat stress and exacerbate existing health conditions.

1 additional hot day per year projected across the region between 2050 and 2090. However, increases of 5.3 additional hot days per year under a medium-emissions scenario and 8.9 additional hot days per year under a high-emissions scenario are projected during the same period.

Changes to the number of hot days per year will vary significantly across the region (Figure 4). The north-western area of the region near Young and Goulburn is projected to experience pronounced increases in the number of hot days. By 2090, Young is projected to experience 13.1 additional hot days per year under a low-emissions scenario, 26.9 under a medium-emissions scenario and 39.9 under a high-emissions scenario. A medium-emissions scenario is projected to more than double Young's baseline period average of 16.7 hot days per year, while a high-emissions scenario is projected to nearly triple Young's baseline average. Comparatively, coastal areas will experience a lower increase due to the moderating influence of the ocean. Batemans Bay's baseline period average is 1 hot day per year. By 2090, Batemans Bay is projected to experience an additional 1.5 hot days per year under a low-emissions scenario, 2.7 under a medium-emissions scenario and 4.2 under a high-emissions scenario.

Table 3 and Figure 3 provide more information on how the projections differ across the 3 scenarios, and Figure 4 provides information on regional differences by 2090 across the 3 scenarios.



Hot days

Table 3. Projected increase in average annual number of hot days – South East and Tablelands

2050


Low-emissions	Medium-emissions	High-emissions
3.9 days (0.7 to 7.3 days)	5.0 days (2.0 to 7.9 days)	7.0 days (1.0 to 15.0 days)


2090

Low-emissions	Medium-emissions	High-emissions
4.7 days (1.2 to 11.0 days)	10.3 days (4.1 to 19.2 days)	15.9 days (7.0 to 30.9 days)

The bold number is the ensemble average for the period. Underneath the average is the ensemble range. Hot day increases are additional to the historical model baseline of 4.4 hot days.


Figure 3. Historical and projected average annual number of hot days – South East and Tablelands





Hot days

Figure 4. Projected change in annual number of hot days by 2090 for the South East and Tablelands





Cold nights

Cold nights are important for biodiversity in higher-elevation areas and the viability of important plant species, including some temperate fruits.

Cold nights will decrease

Cold nights are those where the minimum temperature drops below 2°C. These are important for the viability of some important plant species. For example, some common temperate fruit species require sufficiently cold winters to produce flower buds.

Cold nights in the South East and Tablelands region generally occur for most inland areas of the region, particularly in the Snowy Mountains. During the baseline period, areas of higher elevation in the Snowy Mountains, including regional towns such as Cooma, had on average more than 140 cold nights per year. Lower elevation inland areas, such as Goulburn, Queanbeyan and Young, had on average 70–90 cold nights per year. Areas along the coast rarely experience cold nights, with locations such as Batemans Bay and Eden having 1 cold night per year.

Projections

Across the South East and Tablelands, the average number of cold nights per year will decrease throughout this century (Figure 5).

The number of cold nights will decrease for the region by 2050 for all emissions scenarios, with an even greater decrease by 2090 under a medium-emissions scenario and a high-emissions scenario (Table 4).

Under a high-emissions scenario, alpine areas of the region could experience half the number of cold nights per year by 2090.

Under a low-emissions scenario, the number of cold nights across the South East and Tablelands could reduce by less than 25% by 2090.

The number of cold nights is projected to decrease across autumn, winter and spring, with the largest decreases in winter.

Under a low-emissions scenario, there is a small decrease of 3.4 fewer cold nights per year projected across the region between 2050 and 2090 (Table 4). However, decreases of 16.1 fewer cold nights per year under a medium-emissions scenario and 27.2 fewer cold nights per year under a high-emissions scenario are projected during the same period.

Cold nights will decrease across most of the region, particularly in the higher elevation areas in the south-west (Figure 6). Coastal areas will not experience any changes, as they rarely experience cold nights. The greatest decreases are projected to occur for the Snowy Mountains including Jindabyne and Cooma. By 2090, Cooma is projected to have 22.4 fewer cold nights per year under a low-emissions scenario, 47.2 under a medium-emissions scenario and 69.9 under a high emissions scenario. A medium-emissions scenario is projected to reduce Cooma's 145.1 cold nights per year baseline period average by more than 30%, while a high-emissions scenario is projected to reduce the baseline average by more than 45%.

Table 4 and Figure 5 provide more information on how the projections differ across the 3 scenarios, and Figure 6 provides information on regional differences by 2090 across the 3 scenarios.



Cold nights

Table 4. Projected decrease in average annual number of cold nights – South East and Tablelands

2050

Low-emissions	Medium-emissions	High-emissions
16.4 days (7.9 to 24.0 days)	23.2 days (13.5 to 29.6 days)	28.0 days (15.9 to 36.1 days)

2090

Low-emissions	Medium-emissions	High-emissions
19.8 days (11.9 to 26.0 days)	39.3 days (27.7 to 52.4 days)	55.2 days (43.1 to 69.3 days)

The bold number is the ensemble average for the period. Underneath the average is the ensemble range. Cold night decreases are relative to the historical model baseline of 88.9 cold nights.

Figure 5. Historical and projected change in annual number of cold nights – South East and Tablelands







Figure 6. Projected change in annual number of cold nights by 2090 for the South East and Tablelands





Decreased cold nights


Natural snow depth in alpine areas has declined by over one-third since the 1950s.⁶ Years with persistent heavy snow cover have become rare.⁶ Further reductions in natural snow depth are likely to cause a decline in recreational opportunities, with impacts to local economies dependent on snow-based tourism.

Other significant impacts from climate change can be expected on alpine biodiversity. Critically endangered snowpatch plant communities that depend on long-lasting snow cover may decline in extent as snow cover decreases and competition from other species increases.⁷ The endangered mountain pygmy possum requires snow cover for hibernation and protection from predators. Reduced snow cover from climate change may further restrict available habitat for the mountain pygmy possum.⁸

These ecosystems have minimal adaptive capacity to a decrease in cold nights and subsequent loss of snow cover, as they already occupy the highest elevations and cannot migrate to other areas.^{9,10}



Significant impacts on alpine biodiversity can be expected from reduced snow cover, such as reducing available habitat for the endangered mountain pygmy possum.



Under a low-emissions scenario, the number of cold nights across NSW could reduce by less than 35% by 2090.



Continued decline in snow depth is likely to have significant impacts on the Snowy Mountains Scheme that generates hydro-electric power and provides water for irrigation, which is highly reliant on spring snowmelt and rainfall.⁶





Rainfall

Rainfall is projected to remain variable

Climate change will influence rainfall patterns and the total amount of rainfall that NSW receives. These changes may have widespread impacts on water security, agricultural productivity and native species' reproductive cycles. For example, eucalypt woodlands could struggle to cope with drier conditions.

Modelling rainfall is more difficult than modelling temperature due to the complexities of the weather systems that generate rain. NARcliM projections capture a range of plausible climate futures under the 3 emissions scenarios, including wet and dry outcomes. This means that rainfall is inherently more variable in the NARcliM projections than temperature, and the full range of rainfall projections should be taken into account. This can be explored further on the [AdaptNSW Interactive Map](#).

This snapshot provides data on average rainfall change and does not provide data on rainfall extremes or the impacts of climate change on flooding.

Observed rainfall across the South East and Tablelands region averages about 720 mm.⁴ Rainfall is highest in the Snowy Mountains. The coast and the Southern Highlands also experiences high rainfall, with the area around Goulburn and the Monaro experiencing relatively lower rainfall. Rainfall is nearly uniformly distributed throughout the year, with slightly more rain in summer and autumn. The driest year on record was 1982, which had an average rainfall of only 370 mm. A notably dry year was also experienced in 2019, with approximately 460 mm of rainfall across the region.⁴

A decrease in average winter rainfall of approximately 24–27% by 2090 is projected for the South East and Tablelands under medium-emissions and high-emissions scenarios.

Projections

Annual average rainfall in the region is projected to remain variable throughout this century (Figure 7). By 2090, on average, annual rainfall is projected to decrease by 10% under a low-emissions scenario, by 18% under a medium-emissions scenario and by 12% under a high-emissions scenario (Table 5).

On average, autumn rainfall is projected to change by 10% or less across the region by 2090 under all emissions scenarios. On average, summer rainfall is projected to change by 15% or less across the region by 2090 under all emissions scenarios.

By 2090, on average, winter rainfall is projected to decrease across the region by 8% under a low-emissions scenario, by 28% under a medium-emissions scenario and by 24% under a high-emissions scenario (Table 5). Areas near the Great Dividing Range and in the Southern Highlands are projected to experience the greatest decreases in average winter rainfall. By 2090, for average winter rainfall, Moss Vale is projected to experience a decrease of 23% under a low-emissions scenario, 39% under a medium-emissions scenario and approximately 41% under a high-emissions scenario.

By 2090, on average, spring rainfall is projected to decrease across the region by 12% under a low-emissions scenario, by 19% under a medium-emissions scenario and by 18% under a high-emissions scenario. Areas in the west and north-west of the region including the Snowy Mountains are projected to experience the greatest decreases in average spring rainfall.

Refer to the [Interactive Map](#) for further seasonal information.

Table 5 and Figure 7 provide more information on how the projections differ across the 3 scenarios, and Figures 8 to 12 provides information on regional differences by 2090 across the 3 scenarios by season.



Table 5. Projected change to average rainfall – South East and Tablelands

	Low-emissions	Medium-emissions	High-emissions
Annual	-7.7% (-14.9% to +10.0%)	-7.5% (-20.9% to +4.0%)	-14.4% (-31.0% to -0.5%)
Summer	-7.6% (-22.2% to +22.2%)	-8.0% (-22.9% to +21.0%)	-15.9% (-33.6% to +15.6%)
Autumn	-4.8% (-19.7% to +13.8%)	-0.5% (-26.2% to +36.2%)	-9.6% (-36.6% to +16.1%)
Winter	-9.7% (-22.8% to +32.6%)	-12.1% (-25.8% to +18.9%)	-17.2% (-35.1% to +8.3%)
Spring	-8.7% (-25.2% to +30.4%)	-9.1% (-27.7% to +13.9%)	-14.1% (-23.3% to -1.8%)

	Low-emissions	Medium-emissions	High-emissions
Annual	-9.9% (-22.3% to +11.3%)	-18.2% (-30.8% to +2.1%)	-11.6% (-32.9% to +10.5%)
Summer	-11.9% (-31.1% to +31.9%)	-15.3% (-25.4% to +36.5%)	-4.2% (-38.1% to +40.9%)
Autumn	-6.7% (-24.5% to +13.1%)	-10.3% (-32.5% to +10.2%)	-1.8% (-20.8% to +32.1%)
Winter	-8.3% (-28.3% to +19.8%)	-28.3% (-41.2% to +13.3%)	-23.7% (-40.3% to +13.1%)
Spring	-12.3% (-25.3% to +3.7%)	-19.4% (-31.9% to +6.0%)	-18.3% (-38.4% to -3.5%)

The bold number is the ensemble average for the period. Underneath the average is the ensemble range. Percentages changes in annual average rainfall are relative to the historical model baseline of 743 mm. Average summer rainfall is relative to a baseline of 222 mm, average autumn rainfall is relative to a baseline of 170 mm, average winter rainfall is relative to a baseline of 175 mm and average spring rainfall is relative to a baseline of 175 mm.

Figure 7. Historical and projected change in average rainfall – South East and Tablelands






Figure 8. Projected change to average annual rainfall by 2090 for the South East and Tablelands






Figure 9. Projected change to average summer rainfall by 2090 for the South East and Tablelands






Figure 10. Projected change to average autumn rainfall by 2090 for the South East and Tablelands






Figure 11. Projected change to average winter rainfall by 2090 for the South East and Tablelands







Figure 12. Projected change to average spring rainfall by 2090 for the South East and Tablelands





Severe fire weather

Severe fire weather will increase

The Forest Fire Danger Index (FFDI) represents an estimate of fire weather risk. The FFDI is calculated from temperature, relative humidity and wind speed, as well as an index representing fuel dryness.

Severe fire weather (FFDI greater than 50) is most likely in summer and spring. Fire weather was the strongest determining factor of house loss during the Black Summer bushfires.¹²

The number of severe fire danger days observed for the South East and Tablelands region is 0.5 days per year on average.⁴ The number of severe fire danger days is generally low across the region, with relatively more severe fire danger days in northern inland areas of the region such as Goulburn and Young. The record number of severe fire danger days in a year was 2019 with approximately 3.7 days on average across the region, including 3 days recorded at the Bega weather station and 5 days recorded at the Cooma station.⁴

FFDI was monitored by weather stations across NSW until the introduction in 2022 of the Australian Fire Danger Rating System. FFDI is used in this snapshot as it can be simulated using the NARClIM projections, whereas data used by the Australian Fire Danger Rating System currently cannot. FFDI also provides a long history of data and gives context to the NARClIM projections.

Projections

Across the South East and Tablelands, the average number of severe fire weather days per year will increase throughout this century (Figure 13).

The average number of severe fire weather days will increase for the South East and Tablelands region by 2050 under all emissions scenarios, with an even greater increase projected by 2090 under a medium-emissions scenario and a high-emissions scenario (Table 6). The number of severe fire weather days is projected to increase during spring and summer, with the largest increase in summer.

3x

Under a high-emissions scenario, the number of severe fire weather days per year could triple across the South East and Tablelands by 2090.

Fire weather was the strongest determining factor of house loss during the Black Summer bushfires.¹²

Increases to severe fire weather days will vary across the region (Figure 14). The greatest increases are projected to occur in inland areas including Young, although some coastal areas will also see increases. By 2090, Young is projected to experience 2.0 additional severe fire weather days per year under a low-emissions scenario, 4.5 additional severe fire weather days under a medium-emissions scenario and 5.9 additional severe fire weather days per year under a high-emissions scenario. Both a medium-emissions scenario and a high-emissions scenario are projected to more than double Young's baseline period average of 4.0 severe fire weather days per year. Comparatively, on the coast, Batemans Bay's baseline period average is 0.8 severe fire weather days per year. By 2090, Batemans Bay is projected to experience 0.4 additional severe fire weather days per year under a low-emissions scenario, 0.9 additional severe fire weather days under a medium-emissions scenario and 1.2 additional severe fire weather days per year under a high-emissions scenario.

Table 6 and Figure 13 provide more information on how the projections differ across the 3 scenarios, and Figure 14 provides information on regional differences by 2090 across the 3 scenarios



Table 6. Projected increase in average annual number of severe fire weather days – South East and Tablelands

2050

Low-emissions	Medium-emissions	High-emissions
0.5 days (–0.6 to 1.4 days)	0.6 days (0.0 to 1.6 days)	1.2 days (0.2 to 2.5 days)

2090

Low-emissions	Medium-emissions	High-emissions
0.7 days (–0.5 to 2.1 days)	1.7 days (0.4 to 4.0 days)	2.4 days (0.3 to 5.5 days)

The bold number is the ensemble average for the period. Underneath the average is the ensemble range. Severe fire weather increases are additional to the historical model baseline of 1.2 severe fire weather days.

Figure 13. Historical and projected change in annual number of severe fire weather days – South East and Tablelands







Figure 14. Projected change to average annual number of severe fire weather days by 2090 for the South East and Tablelands





Bushfires

The 2019–20 bushfire season caused extensive damage to communities, infrastructure and natural ecosystems in the South East and Tablelands region. Over 1 million hectares of the region were burnt and 1,135 homes were destroyed.¹³ Large areas of bushland experienced extreme fire severity, including the Beowa, Deua, Wadbilliga and Kosciuszko national parks. Over 27,000 hectares or 21% of NSW alpine vegetation was burnt, adding to extensive areas of alpine vegetation still recovering from severe fires that occurred in 2002–03.¹⁴

The South East and Tablelands region, with its isolated coastal towns near bushland, is highly vulnerable to the impacts of increasing number of days of severe fire weather.

Climate change is also expected to reduce the interval between fires, increase fire intensity, and shorten the window for safe fire management activities.¹⁵ For communities on the bushland-urban interface, the increased fire occurrence heightens risks to people, homes and infrastructure.⁹



North-western areas of the region such as Goulburn and Young will experience greater increases in the number of severe fire weather days.



Although bushfires are a natural part of the Australian environment, climate change is increasing the frequency, duration, extent and intensity of ‘fire weather’ in south-east Australia.^{15,16}



Severe fire danger days, which create the underlying conditions for large-scale bushfires, are expected to become more common in the future, particularly under a high-emissions scenario.





Climate action and further information

Climate action

The NARcliM projections for the low-, medium- and high-emissions scenarios highlight the stark difference in climate change impacts that will be experienced under each scenario. The differences provide a reminder of the required action across the world to reduce emissions, and specifically within NSW to meet our legislated Net Zero by 2050 emissions reduction targets. This is our best chance at ensuring the future projections under the high-emissions scenario are avoided. The NARcliM projections highlight the importance of taking action to adapt to the impacts of climate change. For more resources on reducing emissions and adapting to the impacts of climate change, visit Adapt NSW.

Additional resources

- For information on other climate change impacts, including sea level rise, visit [AdaptNSW](#)
- [Climate change resources for local government on AdaptNSW](#)
- Generate detailed climate information based on your local government area using [SEED](#)
- [Climate Data Portal](#)
- [NARcliM case studies](#)
- [Climate risk ready NSW guide](#)
- [Local government climate change action in NSW: a guide to leadership](#)

Further information

NARcliM projections are delivered with support from: the ACT, South Australian, Victorian and Western Australian governments; National Computational Infrastructure; Murdoch University; and the University of New South Wales. Detailed information on the methodology and application of the projections can be found on the AdaptNSW website. Climate change information in this snapshot is delivered as part of the NSW Government's commitment to 'Publish regularly updated and improved local level climate change projections' under Action 3 of the [NSW climate change adaptation strategy](#).

References

1. O'Neill et al. 2017, The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Global Environmental Change*, 42, 169–180.
2. Riahi et al. 2017, The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*, 42, 153–168.
3. Long-term temperature record – webpage.
4. About Australian Gridded Climate Data maps and grids – webpage.
5. ABS 2016, Census of Population and Housing, TableBuilder.
6. Coates et al. 2014, Exploring 167 years of vulnerability: An examination of extreme heat events in Australia 1844–2010. *Environmental Science & Policy*, 42, 33–44.
7. Varghese et al. 2018, Are workers at risk of occupational injuries due to heat exposure? A comprehensive literature review, *Safety Science*, 110, 380–392.8
8. Beggs et al. 2021, The 2021 report of the MJA-Lancet Countdown on health and climate change: Australia increasingly out on a limb. *The Medical Journal of Australia* 215 (9), 390.
9. Green & Pickering 2002, A scenario for mammal and bird diversity in the Snowy Mountains of Australia in relation to climate change. In *Mountain biodiversity* (pp. 239–247). Routledge.
10. Pittock et al. 2003, Climate change: an Australian guide to the science and potential impacts. Australian Greenhouse Office.
11. Snowy Hydro – webpage
12. Price et al. 2020, Probability of house destruction. Theme 3A. People and Property Impacts, Bushfire Risk Management Research Hub for the NSW Bushfire Inquiry 2020.
13. Owens & O'Kane 2020, Final report of the NSW Bushfire Inquiry, Department of Premier and Cabinet, Sydney.
14. DPIE 2020, NSW fire and the environment 2019–20 summary, Department of Planning, Industry and Environment, Sydney.
15. Hennessy et al. 2007, Australia and New Zealand. In *Climate Change 2007: impacts, adaptation and vulnerability*. Cambridge University Press (CUP), 2007, 507–540.
16. Van Oldenborgh et al. 2021, Attribution of the Australian bushfire risk to anthropogenic climate change. *Natural Hazards and Earth System Sciences Discussions*, 2020, 1–46
17. Dowdy et al. 2019, Future changes in extreme weather and pyroconvection risk factors for Australian wildfires, *Scientific Reports*, 9, 10073

Photo credits:

Cover page: Snow-covered mountains in Kosciuszko National Park, leelakajonkij/iStock; Photograph of dry alpine trees above the timberline in the Snowy Mountains in New South Wales, Phillip Wittke/iStock

Contents:

p.2: The Dark Emu Overhead, Alan Dyer
p.3: Revegetation in Australia, MarkPlovesan/iStock
p.7: Alpine vegetation, Kosciuszko Summit, Kosciuszko National Park. John Spencer/DCCEEW; Aerial view overlooking Central Tilba on the State's South Coast. *Destination NSW*
p.9: Surfer coming out of the ocean at sunrise, Pat Suraseang/*Destination NSW*; Navigate Expeditions, Pambula, SafariGlobal/*Destination NSW*
p.12: Hot day, Serg64/Shutterstock
p.15: Frost on the farm, MacierPhotography/iStock
p.18: Thredbo Resort. Ben Kelly/*Destination NSW*
p.27: Rainforest, New South Wales, Zaharov/iStock
p.30: NPWS fire personnel hosing flames. Phillip Tattersall/DCCEEW
p.31: Putting in a remote Rako line fire control, Michael Jarman/DCCEEW

© 2025 State of NSW

With the exception of photographs, the State of NSW and Department of Climate Change, Energy, the Environment and Water (DCCEEW) are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required for the reproduction of photographs. DCCEEW has compiled this report in good faith, exercising all due care and attention. No representation is made about the accuracy, completeness or suitability of the information in this publication for any particular purpose. DCCEEW shall not be liable for any damage which may occur to any person or organisation taking action or not on the basis of this publication. Readers should seek appropriate advice when applying the information to their specific needs. All content in this publication is owned by DCCEEW and is protected by Crown Copyright, unless credited otherwise. It is licensed under the Creative Commons Attribution 4.0 International (CC BY 4.0), subject to the exemptions contained in the licence. The legal code for the licence is available at Creative Commons. DCCEEW asserts the right to be attributed as author of the original material in the following manner: © State of New South Wales and Department of Climate Change, Energy, the Environment and Water 2024.